



Genesee County Health Department

Mark Valacak, M.P.H., Health Officer
Gary K. Johnson, M.D., M.P.H., Medical Director

FOR IMMEDIATE RELEASE

DATE: November 30, 2012

END DATE: December 24, 2012

CONTACT: Suzanne Cupal, MPH
Epidemiologist

PHONE: (810) 768-7970

RE: Viral Gastroenteritis

Genesee County is experiencing an increase in gastroenteritis cases. The viruses that cause gastroenteritis, such as norovirus, can be easily spread by food, person-to-person contact, or contaminated surfaces or objects. The Genesee Health Department would like to make sure that everyone stays healthy this holiday season by providing the following information.

Norovirus causes an illness which typically includes nausea, vomiting, diarrhea and stomach cramping. Symptoms may also include low-grade fever, headache, fatigue and muscle aches. Symptoms last between 12 and 60 hours. In most cases, ill persons recover on their own. The very young and the elderly are at higher risk for dehydration. Those with severe diarrhea should drink plenty of fluids.

Norovirus is very contagious and can be easily spread from person-to-person, especially in group settings. Individuals can reduce their risk of becoming infected by frequent hand washing using soap and warm water for a minimum of 20 seconds (alcohol based hand sanitizers do **not** prevent the spread of norovirus). Preventing contamination of food, drinks and surfaces is critical to preventing the spread of norovirus. Anyone with norovirus should not prepare or serve food

1 of 2

Better Life Through Better Health

Floyd J. McCree Courts & Human Services Building ♦ 630 S. Saginaw Street, Ste. 4 ♦ Flint, Michigan 48502-1540

Burton Branch ♦ G-3373 S. Saginaw Street ♦ Burton, Michigan 48529

Main Phone 810-257-3612 ♦ Visit us at: www.gchd.us

for others for 3 days after symptoms have stopped. Spread of norovirus can be prevented by disinfecting contaminated surfaces with household chlorine bleach-based cleaners and prompt washing of contaminated articles of clothing. Since the virus is passed in vomit and stool, children should not go to daycare or school while they have diarrhea or vomiting. Persons who work in nursing homes, take care of patients or handle food should stay out of work for 2 to 3 days after symptoms end.

Norovirus is known incorrectly as the “stomach flu.” Norovirus is **not** related to the flu (influenza) which is a respiratory illness caused by a different virus. Attached is a Center for Disease Control and Prevention Fact Sheet regarding norovirus. Additional information about can be found at www.gchd.us.

Norovirus Illness: Key Facts

Norovirus—the stomach bug

Norovirus is a highly contagious virus. Norovirus infection causes gastroenteritis (inflammation of the stomach and intestines). This leads to diarrhea, vomiting, and stomach pain.

Norovirus illness is often called by other names, such as food poisoning and stomach flu. It is true that noroviruses can cause food poisoning. But, other germs and chemicals can also cause food poisoning. Norovirus illness is not related to the flu (influenza), which is a respiratory illness caused by influenza virus.



Anyone can get norovirus illness

- Norovirus is the most common cause of acute gastroenteritis in the United States.
- Each year, norovirus causes about 21 million cases of acute gastroenteritis in this country.
- Many different types of norovirus exist, so you can get infected and sick many times in your life.

Norovirus illness can be serious

- Norovirus illness can make you feel extremely sick with diarrhea and vomiting many times a day.
- Some people may get severely dehydrated, especially young children, the elderly, and people with other illnesses.
- Each year, norovirus causes about 70,000 hospitalizations and 800 deaths, mostly in young children and the elderly.

Norovirus spreads very easily and quickly

- It only takes a very small amount of norovirus particles (fewer than 100) to make you sick.
- People with norovirus illness shed billions of virus particles in their stool and vomit and can easily infect others.
- You are most contagious when you are sick with norovirus illness and during the first 3 days after you recover.
- Norovirus can spread quickly in closed places like daycare centers, nursing homes, schools, and cruise ships.
- Norovirus can stay on objects and surfaces and still infect people after days or weeks.
- Norovirus can survive some disinfectants, making it hard to get rid of.

Norovirus can spread in many ways

Norovirus can spread to others by—

- eating food or drinking liquids that are contaminated with norovirus,
- touching surfaces or objects that have norovirus on them then putting your fingers in your mouth, and
- having close personal contact with an infected person, for example, caring for or sharing food, drinks, or eating utensils with an infected person.

There's no vaccine to prevent norovirus infection and no drug to treat it

- Antibiotics will not help with norovirus illness because antibiotics do not work on viruses.
- When you have norovirus illness, drink plenty of liquids to replace fluid loss and prevent dehydration.
- If you or someone you are caring for is dehydrated, call a doctor.

National Center for Immunization and Respiratory Diseases
Division of Viral Diseases

CS234745-A





What is the Right Way to Wash Your Hands?

1. Wet your hands with clean, running water (warm or cold) and apply soap.
2. Rub your hands together to make a lather and scrub them well; be sure to scrub the backs of your hands, between your fingers, and under your nails.
3. Continue rubbing your hands for at least 20 seconds. Need a timer? Hum the "Happy Birthday" song from beginning to end twice.
4. Rinse your hands well under running water.
5. Dry your hands using a clean towel or air dry them.

See Handwashing: Clean Hands Saves Lives (www.cdc.gov/handwashing/)

5 Tips to Prevent Norovirus From Spreading

1. Practice proper hand hygiene

Always wash your hands carefully with soap and water—

- after using the toilet and changing diapers, and
- before eating, preparing, or handling food.

Alcohol-based hand sanitizers can be used in addition to hand washing. But, they should not be used as a substitute for washing with soap and water.

2. Wash fruits and vegetables and cook seafood thoroughly

Carefully wash fruits and vegetables before preparing and eating them.

Cook oysters and other shellfish thoroughly before eating them.

Be aware that noroviruses are relatively resistant. They can survive temperatures as high as 140°F and quick steaming processes that are often used for cooking shellfish.

Food that might be contaminated with norovirus should be thrown out.

Keep sick infants and children out of areas where food is being handled and prepared.

3. When you are sick, do not prepare food or care for others

You should not prepare food for others or provide healthcare while you are sick and for at least 2 to 3 days after you recover. This also applies to sick workers in schools, daycares, and other places where they may expose people to norovirus.

4. Clean and disinfect contaminated surfaces

After throwing up or having diarrhea, immediately clean and disinfect contaminated surfaces. Use a chlorine bleach solution with a concentration of 1000–5000 ppm (5–25 tablespoons of household bleach [5.25%] per gallon of water) or other disinfectant registered as effective against norovirus by the Environmental Protection Agency (EPA).

5. Wash laundry thoroughly

Immediately remove and wash clothes or linens that may be contaminated with vomit or stool (feces).

You should—

- handle soiled items carefully without agitating them,
- wear rubber or disposable gloves while handling soiled items and wash your hands after, and
- wash the items with detergent at the maximum available cycle length then machine dry them.



Visit CDC's Norovirus Web site at www.cdc.gov/norovirus for more information.